






HPCMP CREATE™-AV Kestrel Simulations for the DPW-8/AePW-4 Buffet Working Group: Part 1

APA-31: DPW-8/AePW-4: Buffet Working Group – ONERA
OAT15A Test Case
Wednesday, 23 July 2025

HPCMP CREATE™-AV Kestrel



- Multiphysics simulation code supporting US DoD fixed-wing programs of record
 - Full spectrum of flight conditions and missions
 - Full spectrum of aircraft/weapons types
 - Coupled physics: **aerodynamics**, thermochemistry, aero-heating, **structural dynamics**, propulsion, flight controls, 6-DoF motion w/ generalized constraints, separation/contact/collision
 - Easy for users to learn, use, customize, and extend



Problem Description

- **ONERA OAT15A transonic airfoil**



- Mach 0.73, $Re_c = 3 \times 10^6$, $T_\infty = 271$ K
- Experimental C_p and $RMS(C_p)$ available at several α

- **Workshop Problems**

- Case 1a – use a steady solver to solve at $\alpha = \{1.36^\circ, 1.50^\circ, 2.50^\circ, 3.00^\circ, 3.10^\circ, 3.25^\circ, 3.40^\circ, 3.50^\circ, 3.60^\circ, 3.90^\circ\}$
- Case 1b – use unsteady solver to solve at same α
 - Focus on grid level 3


- **Kestrel Setup**

- Use workshop-provided unstructured grids from Cadence
- Look at both Spalart-Allmaras (SA) and Menter Shear Stress Transport (SST) turbulence models
 - Both rotational/curvature correction and QCR 2020 turned on for both models
- Using Kestrel's "FN1+NN" expanded gradient stencil




CASE 1A


Steady Solver Results: SA



Steady Solver Results: SST




Steady Solver Results: C_p Distributions




CASE 1B

Unsteady Solver Time Step Study: $\alpha=3.90^\circ$, SA





- C_L , C_D , C_M all move noticeably as time step changes
- Increasing number of subiterations does not have significant impact on behavior

Unsteady Solver Time Step Study: $\alpha=3.90^\circ$, SST



- C_L , C_D , C_M all move noticeably as time-step changes, but less so than for SA
- Subiterations matter
 - 3 subiterations produce steady-ish behavior, 10 subiterations produce unsteady behavior

Unsteady Solver Time Step Study: $\alpha=3.90^\circ$, C_p



- For SA, no discernable difference between 3 and 10 subiterations
- For SST, 3 subiterations shows similar behavior to steady response, and 10 subiterations shows significantly improved comparison to experimental data

Conclusions

- **Case 1a – Steady Simulations (RANS)**

- Conducted steady runs at all grid levels using both SA and SST.
- Both SA and SST simulations achieved reasonable comparisons to experimental data when comparing mean C_p distribution.
- The SA shock location tended to be further downstream than the SST shock location. The experimental shock location was between the two for lower α and upstream of both at higher α .
- SST showed signs that the flow was unsteady at most grid levels for $\alpha \geq 3.10^\circ$, but SA only demonstrated signs of potential unsteadiness for the L6 grid.

- **Case 1b – Unsteady Simulations (URANS)**

- Conducted initial time step study at $\alpha = 3.90^\circ$ on the L3 grid using both SA and SST.
- For SA, a suitable time step and subiteration count combination has not been found.
- For SST, multiple time steps using 10 subiterations showed reasonable unsteady results when compared to the experimental data.

Acknowledgements

- Material presented here is a product of the Computational Research and Engineering for Acquisition Tools and Environments (CREATE) Program sponsored by the U.S. Department of Defense HPC Modernization Program Office.
- Computations done in support of this effort were done on supercomputing resources provided by the HPCMP at the Navy DoD Supercomputing Resource Center (DSRC) and by the High-End Computing Capability (HECC) at the NASA Advanced Supercomputing (NAS) Facility.

QUESTIONS?